

AD-A106 356

MASSACHUSETTS UNIV AMHERST
AROMATIC CONDUCTING POLYMERS. (U)
OCT 81 J C CHIEN, C P LILLYA
TR-1

F/G 11/9

UNCLASSIFIED

N00014-80-C-0756
NL

For
Date



END
DATE
FILMED
11-81
DTIC

AD A106356

LEVEL

(12)

OFFICE OF NAVAL RESEARCH

Contract N00014-80-C-0756

TECHNICAL REPORT NO. 1

Aromatic Conducting Polymers.

by

James C.W. Chien and C. Peter Lillya

DTIC
ELECTE
OCT 30 1981
S H

University of Massachusetts
Department of Polymer Science and Engineering
Department of Chemistry
Amherst, Massachusetts 01003

October 1981

Reproduction in whole or in part is permitted for
any purpose of the United States Government

* This document has been approved for public release
and sale; its distribution is unlimited

* This statement should also appear in Item 10 of Document Control
Data - DD Form 1473. Copies of form available from cognizant
contract administrator.

UIC FILE COPY

87 9 8 185

Annual Report (October, 1981)

Aromatic Conducting Polymers (Contract #N00014-80-C-0756)

Principal Investigators: James C. W. Chien and C. Peter Lillya

Interests in electrically conducting polymers have grown from polyacetylene to other polymers. One of the objectives of this research project, initiated this year, is in elucidating structure-conductivity relationships for new aromatic polymers.

The following model compounds and polymers have been synthesized: poly(2,5-dimethoxy-phenylene-vinylene), para-trans-trans-distyrylbenzene, meta-trans-trans-distyrylbenzene, trans-poly(phenylene-vinylene) both by Wittig reaction and by NaH synthesis, cis- and trans-poly(phenylene-vinylene) also by both methods, trans-1,4-bis-4-phenylbutadienylbenzene, cis- and trans-1,4-bis-4-phenylbutadienylbenzene, poly-(p-diethynylbenzene) of 4 different molecular weights, poly(m-diethynylbenzene), poly(m-phenylene-vinylene), poly(m- and p-phenylene-vinylene) and trans-stilbene.

Conductivity of the above materials had been obtained before and after AsF₅ doping. The conducting properties are related to: (1) cis, trans structure, (2) molecular weight, (3) method of synthesis, (4) meta and para substitution, and (5) bond alteration.

The formation of radical cations upon AsF₅ doping and subsequent chain extension in these materials are studied by electron paramagnetic resonance for linewidth, g-values, spin concentration and their temperature dependence.

Accession For

NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	

By _____

Distribution/

Availability Codes

Avail and/or	Special
Dist	

A

